



Agricultural Research Division
 University of Nebraska Extension
 Institute of Agriculture and Natural Resources
 University of Nebraska–Lincoln

2015 Beef Cattle Report



Extension is a Division of the Institute of Agriculture and Natural Resources at the University of Nebraska–Lincoln cooperating with the Counties and the United States Department of Agriculture. University of Nebraska–Lincoln Extension educational programs abide with the nondiscrimination policies of the University of Nebraska–Lincoln and the United States Department of Agriculture.

© 2014, The Board of Regents of the University of Nebraska. All rights reserved.

Nutrient Digestibility and Ruminal pH of Finishing Diets Containing Dry Milling Byproducts With and Without Oil Extraction

Melissa L. Jolly-Breithaupt
Adam L. Shreck
Jana L. Harding
Jim C. MacDonald
Terry J. Klopfenstein
Galen E. Erickson¹

Procedure

A 112-day metabolism experiment utilized six ruminally fistulated steers in a 5 x 5 Latin Square design. Treatments were designed as a 2 x 2 + 1 factorial arrangement with steers assigned randomly to one of five treatments (Table 1). Factors consisted of oil concentration (de-oiled or normal) and byproduct type (27% CDS or 40% MDGS) plus a corn-based control. All diets contained (DM basis) a 1:1 blend of dry-rolled and high-moisture corn which was replaced by either CDS or

MDGS, 12% corn silage, and a 5% supplement. All supplements contained Rumensin[®] and tylosin at 345 and 90 mg per steer daily, respectively. The byproducts utilized in this trial were procured from Green Plains, LLC (Central City, Neb).

Steers were housed in individual concrete slatted pens with *ad libitum* access to feed and water. Ingredient samples were taken during the collection period at time of mixing, composited by period, ground through a 1-mm screen, and analyzed for DM, fat, CP, S, and NDF. Fat concentration

Summary

A metabolism trial was conducted to determine the effects of corn oil removal in condensed distillers solubles (CDS) and modified distillers grains plus solubles (MDGS) on nutrient digestibility and ruminal pH. Oil removal had no impact on DM, OM, or NDF digestibility in steers fed CDS or MDGS. However, steers fed de-oiled CDS had a lower fat digestibility than steers fed normal CDS. Average ruminal pH was lower for steers fed de-oiled MDGS than for steers fed normal MDGS, however no difference within CDS was observed.

Introduction

For the last two years, ethanol plants have been removing a portion of corn oil via centrifugation to produce de-oiled distillers byproducts. Previous research has concluded that removal of corn oil by this centrifugation process had limited impact on ADG and F:G when 27% inclusion of CDS or 40% inclusion of MDGS were fed in finishing diets (2013 *Nebraska Beef Cattle Report*, pp. 64–65). No data have been reported on the nutrient digestibility of diets containing de-oiled byproducts. The hypothesis of this trial was that oil removal would improve NDF digestibility. Therefore, the objective of this study was to determine the effects of feeding de-oiled CDS and MDGS on nutrient digestibility and ruminal pH of finishing steers.

Table 1. Diet composition on a DM basis fed to finishing steers.

| Ingredient, % of DM | Control | 27% CDS ¹ | | 40% MDGS ¹ | |
|-------------------------------|---------|----------------------|------------|-----------------------|------------|
| | | De-Oiled | Normal Fat | De-Oiled | Normal Fat |
| DRC ¹ | 41.5 | 28 | 28 | 21.5 | 21.5 |
| HMC ¹ | 41.5 | 28 | 28 | 21.5 | 21.5 |
| MDGS: De-oiled ¹ | — | — | — | 40 | — |
| MDGS: Normal fat ¹ | — | — | — | — | 40 |
| CDS: De-oiled ¹ | — | 27 | — | — | — |
| CDS: Normal fat ¹ | — | — | 27 | — | — |
| Corn silage | 12 | 12 | 12 | 12 | 12 |
| Supplement ^{2,3} | 5 | 5 | 5 | 5 | 5 |
| Analyzed Composition, % | | | | | |
| Fat | 4.01 | 5.17 | 6.99 | 5.93 | 7.16 |
| CP | 12.4 | 14.8 | 13.8 | 19.0 | 18.5 |
| NDF | 13.1 | 10.2 | 11.9 | 19.9 | 22.6 |
| S | 0.14 | 0.30 | 0.25 | 0.28 | 0.27 |

¹CDS = Condensed distillers solubles; MDGS = Modified distillers grains plus solubles; DRC = Dry-rolled corn; HMC = High-moisture corn.

²Formulated to contain 345 mg/steer daily of Rumensin[®] and 90 mg/steer daily of Tylan[®].

³Control supplement contained 1.516% urea.

Table 2. Nutrient Composition of MDGS and CDS¹.

| | De-Oiled CDS ² | Normal CDS ² | De-Oiled MDGS ² | Normal MDGS ² |
|--------|---------------------------|-------------------------|----------------------------|--------------------------|
| Fat, % | 8.7 | 15.4 | 9.2 | 12.3 |
| CP, % | 29.9 | 25.5 | 33.9 | 32.4 |
| S, % | 0.73 | 0.56 | 0.51 | 0.48 |
| NDF, % | 1.9 | 8.2 | 29.7 | 36.4 |

¹All values expressed on a DM basis.

²CDS = Condensed distillers solubles; MDGS = Modified distillers grains plus solubles.

Table 3. Effects of dietary treatment on intake and total tract digestibility of DM, organic matter, fat, and NDF.

| Item | Control | 27% CDS ¹ | | 40% MDGS ¹ | | SEM | P-value | | | |
|------------------------------|--------------------|----------------------|--------------------|-----------------------|--------------------|------|-------------------|------------------|-------------------|---------------------|
| | | De-Oiled | Normal | De-Oiled | Normal | | Int. ² | CDS ³ | MDGS ⁴ | F-Test ⁵ |
| DM | | | | | | | | | | |
| Intake, lb/day | 22.2 ^{bc} | 19.9 ^a | 21.0 ^{ab} | 24.2 ^c | 22.8 ^{bc} | 1.3 | 0.33 | 0.34 | 0.29 | 0.05 |
| Total tract digestibility, % | 81.6 | 81.4 | 83.6 | 82.1 | 80.0 | 1.9 | 0.14 | 0.17 | 0.26 | 0.27 |
| OM | | | | | | | | | | |
| Intake, lb/day | 21.3 ^{bc} | 18.7 ^a | 19.9 ^{ab} | 23.1 ^c | 21.8 ^c | 1.2 | 0.33 | 0.29 | 0.32 | 0.03 |
| Total tract digestibility, % | 82.9 ^{ab} | 84.6 ^{bc} | 86.0 ^c | 83.6 ^{abc} | 81.9 ^a | 1.8 | 0.21 | 0.30 | 0.30 | 0.08 |
| NDF | | | | | | | | | | |
| Intake, lb/day | 2.9 ^b | 1.9 ^a | 2.1 ^c | 4.7 ^d | 5.1 ^e | 0.2 | 0.40 | 0.43 | 0.06 | <0.01 |
| Total tract digestibility, % | 58.0 | 53.6 | 61.0 | 67.0 | 67.0 | 5.5 | 0.38 | 0.17 | 0.99 | 0.12 |
| Fat | | | | | | | | | | |
| Intake, lb/day | 0.90 ^a | 1.02 ^a | 1.46 ^b | 1.46 ^b | 1.64 ^c | 0.08 | 0.07 | <0.01 | 0.05 | <0.01 |
| Total tract digestibility, % | 87.3 ^a | 89.6 ^{ab} | 93.1 ^c | 91.2 ^{bc} | 90.6 ^b | 1.2 | 0.03 | 0.02 | 0.68 | 0.01 |

^{a-c}Means with different superscripts differ ($P < 0.10$).

¹27% CDS = 27% inclusion of condensed distillers solubles; 40% MDGS = 40% inclusion of modified distillers grains plus solubles.

²Int = Interaction P-value for byproduct type and oil concentration.

³CDS = Pair-wise, contrast of de-oiled vs. normal CDS.

⁴MDGS = Pair-wise, contrast of de-oiled vs. normal MDGS.

⁵F-Test = Overall F-test representing variation due to treatment.

was analyzed using the biphasic lipid extraction procedure with NDF analyzed after fat had been extracted.

Period duration was 21 days with a 16 day adaptation phase and 5 day collection period. Beginning on day 10 of each period, titanium dioxide was dosed intraruminally at 0800 and 1600 hours to provide a total of 20 g/day. On day 17 to 21, fecal grab samples were collected three times/day at 0800, 1200, and 1600 hours and composited by steer and period. Fecal samples were analyzed for titanium dioxide to determine nutrient digestibility. Fecal samples were also analyzed for DM, organic matter (OM), fat, and NDF. Ruminant pH was measured continuously from day 17 to 21 with submersible wireless pH probes. Measurements for pH included average ruminal pH, minimum and maximum pH, magnitude of change, variance, and time and area below 5.6.

Digestibility, intake, and ruminal pH data were analyzed using the MIXED procedure of SAS (SAS Institute, Inc., Cary, N.C.). Treatment and period were included in the model as fixed effects while steer was treated as a random effect for all analyses. Pair-wise comparisons of treatments were determined by Fisher's LSD and two pre-planned contrasts were used to evaluate the effect of oil removal when

27% CDS or 40% MDGS were fed. Treatment differences were considered significant at $P < 0.10$.

Results

Dietary fat was 5.17% for 27% de-oiled CDS, 6.99% for 27% normal CDS, 5.93% for 40% de-oiled MDGS, and 7.16% for 40% normal MDGS compared to 4.01% fat for the control treatment. The nutrient analysis of CDS and MDGS are included in Table 2.

Intakes

No byproduct by fat concentration interactions were observed for intakes of DM, OM, or NDF ($P \geq 0.33$; Table 3). There were no differences due to oil removal for both 40% MDGS and 27% CDS on DMI and OM intake ($P \geq 0.29$ and $P \geq 0.29$, respectively). However, cattle fed 40% normal MDGS had greater intakes of NDF compared to cattle fed 40% de-oiled MDGS ($P = 0.06$). A byproduct by fat concentration interaction was observed for fat intake ($P = 0.07$). A greater fat intake difference, due to oil removal, was observed for cattle fed 27% CDS compared to 40% MDGS. This response should be expected as the corn oil was only removed from

the CDS portion of the MDGS production process. When comparing all treatments, cattle fed 27% CDS had the lowest ($P < 0.01$) NDF intakes with cattle fed 40% MDGS having the greatest intakes and the control being intermediate. Cattle fed 27% de-oiled CDS had the lowest ($P \leq 0.05$) DM and OM intakes with cattle fed 40% de-oiled MDGS having the greatest intakes and 27% normal CDS, 40% normal MDGS, and control being intermediate.

Digestibility

No byproduct by fat concentration interactions were observed for DM, OM, and NDF digestibilities ($P \geq 0.14$; Table 3). Oil removal had no impact on DM, OM, and NDF digestibility for either cattle fed 27% CDS or 40% MDGS ($P \geq 0.17$). This contradicts our hypothesis that oil removal would improve nutrient digestibility. A byproduct by fat concentration interaction was observed for fat digestibility ($P = 0.03$). The magnitude of difference between de-oiled and normal was greater for cattle fed CDS than for MDGS. Cattle fed 27% normal CDS had a greater fat digestibility compared to 27% de-oiled CDS ($P = 0.02$), whereas no difference

(Continued on next page)

Table 4. Effects of dietary treatment on ruminal pH with steers fed 27% CDS and 40% MDGS with (de-oiled) or without (normal) a portion of oil removed.

| Item | Control | 27% CDS ¹ | | 40% MDGS ¹ | | SEM | P-value | | | |
|--------------------------------------|-------------------|----------------------|-------------------|-----------------------|-------------------|-------|-------------------|------------------|-------------------|---------------------|
| | | De-Oiled | Normal | De-Oiled | Normal | | Int. ² | CDS ³ | MDGS ⁴ | F-Test ⁵ |
| Average pH | 5.40 ^a | 5.39 ^a | 5.36 ^a | 5.54 ^a | 5.72 ^b | 0.09 | 0.14 | 0.85 | 0.09 | 0.02 |
| Maximum pH | 6.05 | 6.15 | 6.02 | 6.19 | 6.38 | 0.11 | 0.21 | 0.43 | 0.24 | 0.21 |
| Minimum pH | 4.99 | 4.95 | 4.98 | 4.93 | 5.09 | 0.09 | 0.16 | 0.78 | 0.18 | 0.68 |
| pH magnitude | 1.08 | 1.15 | 1.02 | 1.30 | 1.31 | 0.13 | 0.60 | 0.51 | 0.92 | 0.44 |
| pH variance ⁶ | 0.072 | 0.101 | 0.065 | 0.074 | 0.131 | 0.023 | 0.33 | 0.28 | 0.10 | 0.26 |
| Time < 5.6, minutes/day ⁷ | 708 | 748 | 1080 | 733 | 769 | 104 | 0.37 | 0.04 | 0.81 | 0.12 |
| Area < 5.6, minutes/day ⁸ | 275 | 312 | 450 | 212 | 302 | 77 | 0.20 | 0.23 | 0.42 | 0.32 |

^{a-c}Means with different superscripts differ ($P < 0.10$).

¹27% CDS = 27% inclusion of condensed distillers solubles; 40% MDGS = 40% inclusion of modified distillers grains plus solubles.

²Int = Interaction P-value for byproduct type and oil concentration.

³CDS = Pairwise, contrast of de-oiled vs. normal CDS.

⁴MDGS = Pairwise, contrast of de-oiled vs. normal MDGS.

⁵F-Test = Overall F-test representing variation due to treatment.

⁶Variance of daily ruminal pH.

⁷Time < 5.6 = minutes that ruminal pH was below 5.6.

⁸Area < 5.6 = ruminal pH units below 5.6 by minute.

was observed between 40% de-oiled and normal MDGS ($P = 0.68$). When comparing all treatments, no differences were observed for DM digestibility ($P = 0.27$) or NDF digestibility ($P = 0.12$). However, steers fed 27% normal CDS had the greatest ($P = 0.08$) OM digestibility, while steers fed 40% normal MDGS had the lowest OM digestibility. Steers fed 27% normal CDS had the greatest ($P = 0.01$) fat digestibility, while control had the lowest.

Ruminal pH

No byproduct by fat concentration interactions were observed for all ruminal pH variables ($P \geq 0.14$; Table 4). Oil removal had no impact

on ruminal pH in steers fed 27% CDS ($P \geq 0.23$) except for time spent below a pH of 5.6 ($P = 0.04$). Steers fed 27% normal CDS spent more time with a ruminal pH below 5.6 than steers fed 27% de-oiled CDS. Oil removal had an effect on average ruminal pH ($P = 0.09$) and pH variance ($P = 0.10$) in steers fed 40% MDGS. Average ruminal pH and variance were lower for steers fed 40% de-oiled MDGS compared to 40% normal MDGS. A treatment effect was observed for average ruminal pH ($P < 0.02$) with steers fed 40% normal MDGS having a greater ruminal pH than steers fed control, 27% de-oiled or normal CDS, and 40% de-oiled MDGS.

These data indicate that oil removal via centrifugation in dry

milling byproducts has limited impact on digestibility in finishing cattle diets. These findings do not support our hypothesis of improved digestibilities in cattle fed de-oiled byproducts; however, it supports the findings as to why there have been little differences observed in finishing performance between de-oiled and normal byproducts.

¹Melissa L. Jolly-Breithaupt, graduate student; Adam L. Shreck, research technician; Jana L. Harding, research technician; Jim C. MacDonald, associate professor; Terry J. Klopfenstein, professor; Galen E. Erickson, professor, University of Nebraska–Lincoln Department of Animal Science, Lincoln, Neb.